

Existence and Uniqueness of Periodic Solutions for (2n + 1)th-Order Differential Equations

Cong Fuzhong, Huang Qingdao, and Shi Shaoyun

metadata, citation and similar papers at core.ac.uk

People's Republic of China

Submitted by Jack K. Hale

Received October 6, 1997

Applying the Schauder fixed-point theorem and the continuation theorem of Mawhin, we give the existence and uniqueness results of periodic solutions for a class of (2n + 1)th-order nonlinear ordinary differential equations. © 2000 Academic Press

Key Words: (2n + 1)th-order differential equation; periodic solution; existence and uniqueness; Schauder fixed-point theorem; continuation theorem of Mawhin.

We consider the (2n + 1)th-order nonlinear ordinary differential equation

$$x^{(2n+1)} + \sum_{i=0}^{n-1} c_i x^{(2i+1)} + f(t, x) = 0, \quad (1)$$

where $f: R \times R \rightarrow R$ is continuous and 2π -periodic with respect to t and c_i , $i = 1, 2, \dots, n-1$, are constants.

During the past 20 years, there has been a great amount of work in periodic solutions for the high-order Duffing equation

$$y^{(2n)} + \sum_{i=1}^{n-1} c_i y^{(2i)} + (-1)^{(n+1)} f(t, y) = 0.$$

Among previous results [1–4], one fundamental nonresonance condition

$$\begin{aligned} 0 &\leq N^{2n} + \sum_{j=1}^{n-1} (-1)^{j-n} c_j N^{2j} \\ &< \lambda \leq f_y(t, y) \leq \mu \\ &< (N+1)^{2n} + \sum_{j=1}^{n-1} (-1)^{j-n} c_j (N+1)^{2j}, \end{aligned}$$

where N is a nonnegative integer, is required. However, the study of odd-order differential equation is rare.

For $c_i = 0$, $i = 1, 2, \dots, n-1$, and f satisfies

$$M \geq f_x(t, x) \geq \varepsilon_0 > 0 \quad (\text{or } -M \leq f_x(t, x) \leq -\varepsilon_0 < 0),$$

Li Yong and Wang Huaizhong [4] proved that (1) has a unique 2π -periodic solution.

In this note, using the Schauder fixed-point theorem and the continuation theorem of Mawhin, we give results on the existence and uniqueness of the 2π -periodic solution to (1). Assume that the following conditions hold:

(H1) f_x is continuous and there are positive constants m and M such that

$$m \leq |f_x(t, x)| \leq M \quad \text{for all } (t, x). \quad (2)$$

(H2) The linear equation

$$x^{(2n+1)} + \sum_{i=0}^{n-1} c_i x^{(2i+1)} = 0 \quad (3)$$

has only the constants for its 2π -periodic solutions.

(H3) $c_j \geq 0$, $j = 0, 1, \dots, n-1$, satisfies the inequality

$$\sum_{j=0}^{n-1} (2\pi)^{2(n-j)} c_j < 1. \quad (4)$$

Our results are the following.

THEOREM A. *Under conditions (H1) and (H2) above, the differential equation (1) has at least one 2π -periodic solution.*

THEOREM B. *Under conditions (H1) and (H3) above, the differential equation (1) has a unique 2π -periodic solution.*

To prove the theorems, we need the following lemma.

LEMMA 1. *Let $x \in C^1(R)$ and let it be a 2π -periodic function; and there exists $t_0 \in R$ such that $x(t_0) = 0$. Then*

$$\int_0^{2\pi} x^2(t) dt \leq 4\pi^2 \int_0^{2\pi} [x'(t)]^2 dt.$$

Employing the Schwarz inequality, the proof of Lemma 1 is easy. Set

$$X = \{y \in C^{2n}(R): y(t) = y(t + 2\pi), \quad \text{for all } t \in R\},$$

with the norm $\|\cdot\|$ defined by $\|y\| = \sum_{i=0}^{2n} \max_{[0, 2\pi]} |y^{(i)}(t)|$, for each $y \in X$.

Rewrite (1) in the following form:

$$x^{(2n+1)} + \sum_{i=0}^{n-1} c_i x^{(2i+1)} + g(t, x)x = -f(t, 0), \quad (5)$$

$$g(t, x) = \int_0^1 f_x(t, \theta x) d\theta.$$

For each $y \in X$, we consider the linear differential equation

$$x^{(2n+1)} + \sum_{i=0}^{n-1} c_i x^{(2i+1)} + g(t, y)x = -f(t, 0). \quad (6)_y$$

LEMMA 2. *There is $M_1 > 0$ such that any 2π -periodic solution $x_y(t)$ of $(6)_y$ satisfies*

$$\|x_y\| \leq M_1, \quad \text{for all } y \in X.$$

Proof. Let $x_y(t)$ be any 2π -periodic solution of $(6)_y$. By Rolle's theorem, we have $\xi_j \in [0, 2\pi]$ satisfying

$$x_y^j(\xi_j) = 0, \quad j = 1, 2, \dots, 2n. \quad (7)$$

Applying Lemma 1,

$$\begin{aligned}
& \left(\int_0^{2\pi} \left(x_y^{(j)} \right)^2 dt \right)^{1/2} \\
& \leq 2\pi \left(\int_0^{2\pi} \left(x_y^{(2j+1)} \right)^2 dt \right)^{1/2} \\
& \leq (2\pi)^{2n-j} \left(\int_0^{2\pi} \left(x_y^{(2n)} \right)^2 dt \right)^{1/2}, \quad j = 1, 2, \dots, 2n-1. \quad (8)
\end{aligned}$$

From (5),

$$m \leq |g(t, x)| \leq M, \quad \text{for all } (t, x). \quad (9)$$

Let $L = \sup_{t \in R} |f(t, 0)|$. Take the product (6)_y with $x_y(t)$ and then integrate over $[0, 2\pi]$. It follows that

$$\begin{aligned}
& \int_0^{2\pi} x_y^{(2n+1)} x_y dt + \sum_{i=0}^{n-1} c_i \int_0^{2\pi} x_y^{(2i+1)} x_y dt + \int_0^{2\pi} g(t, y) x^2 dt \\
& = - \int_0^{2\pi} f(t, 0) x_y dt.
\end{aligned}$$

Noting that $\int_0^{2\pi} x_y^{(2i+1)} x_y dt = 0$ and using (9) and the Schwarz inequality, we derive

$$\left(\int_0^{2\pi} x_y^2 dt \right)^{1/2} \leq \frac{1}{m} \left(\int_0^{2\pi} |f(t, 0)|^2 dt \right)^{1/2} \leq \frac{\sqrt{2\pi}}{m} L. \quad (10)$$

From (6)_y and the Schwarz inequality, using Lemma 1, (10), and $2ab \leq \varepsilon a^2 + b^2/\varepsilon$, we have

$$\begin{aligned}
& \int_0^{2\pi} \left(x_y^{(2n)} \right)^2 dt = - \int_0^{2\pi} \left(x_y^{(2n+1)} \right) \left(x_y^{(2n-1)} \right) dt \\
& = \int_0^{2\pi} \left(\sum_{i=0}^{n-1} c_i x_y^{(2i+1)} x_y^{(2n-1)} \right) dt \\
& \quad + \int_0^{2\pi} g(t, y) x_y x_y^{(2n-1)} dt + \int_0^{2\pi} f(t, 0) x_y^{(2n-1)} dt \\
& \leq \sum_{i=0}^{n-1} c_i \left(\int_0^{2\pi} \left(x_y^{(2i+1)} \right)^2 dt \right)^{1/2} \left(\int_0^{2\pi} \left(x_y^{(2n-1)} \right)^2 dt \right)^{1/2}
\end{aligned}$$

$$\begin{aligned}
& + M \int_0^{2\pi} |x_y x_y^{(2n-1)}| dt + \int_0^{2\pi} L |x_y^{(2n-1)}| dt \\
& \leq \sum_{i=0}^{n-1} c_i (2\pi)^{2(n-i)} \int_0^{2\pi} (x_y^{(2n)})^2 dt + \frac{1}{2} M \varepsilon \int_0^{2\pi} (x_y^{(2n-1)})^2 dt \\
& \quad + \frac{1}{2\varepsilon} M \int_0^{2\pi} x_y^2 dt + \frac{1}{2} \varepsilon \int_0^{2\pi} (x_y^{(2n-1)})^2 dt + \frac{L^2}{2\varepsilon} \cdot 2\pi \\
& \leq \left(2\pi^2 M \varepsilon + 2\pi^2 \varepsilon + \sum_{i=0}^{n-1} c_i (2\pi)^{2(n-i)} \right) \int_0^{2\pi} (x_y^{(2n)})^2 dt \\
& \quad + \frac{\pi L^2}{\varepsilon} \left(1 + \frac{M}{m^2} \right).
\end{aligned}$$

Choose $\varepsilon > 0$ so that

$$1 - \sum_{j=0}^{n-1} (2\pi)^{2(n-j)} c_j - 2\pi^2 \varepsilon - 2\pi^2 M \varepsilon \triangleq A > 0;$$

then we get

$$\int_0^{2\pi} (x_y^{(2n)})^2 dt \leq \frac{\pi L^2}{A \varepsilon} \left(1 + \frac{M}{m^2} \right) \triangleq B^2 L^2. \quad (11)$$

On the basis of (7), (8), (11), and the Schwarz inequality, we obtain

$$\begin{aligned}
\left| x_y^{(j)}(t) \right| &= \left| \int_{\xi_j}^t x_y^{(j+1)} dt \right| \\
&\leq (2\pi)^{1/2} \left(\int_0^{2\pi} (x_y^{(j+1)})^2 dt \right)^{1/2} \\
&\leq (2\pi)^{2n-j+1/2} BL, \\
&\text{for all } t \in [0, 2\pi], \quad j = 1, 2, \dots, 2n-1. \quad (12)
\end{aligned}$$

Also by (6)_y,

$$\int_0^{2\pi} g(t, y) x_y dt = - \int_0^{2\pi} f(t, 0) dt.$$

Using (9), there exists $\xi \in [0, 2\pi]$ satisfying

$$|x_y(\xi)| \leq \frac{1}{2\pi m} \int_0^{2\pi} |f(t, 0)| dt \leq \frac{L}{m}. \quad (13)$$

From (8), (11), and (13),

$$\begin{aligned} |x_y(t)| &\leq |x_y(\xi)| + \left| \int_{\xi}^t x_y(t) dt \right| \\ &\leq \frac{L}{m} + (2\pi)^{2n-1/2} BL \triangleq LD, \quad \text{for all } t \in [0, 2\pi]. \end{aligned} \quad (14)$$

According to (6)_y, for all $t \in [0, 2\pi]$,

$$\begin{aligned} |x_y^{(2n)}(t)| &= \left| \int_{\xi_{2n}}^t x_y^{(2n+1)}(t) dt \right| \\ &\leq \sum_{i=0}^{n-1} c_i \int_0^{2\pi} |x_y^{(2i+1)}| dt + \int_0^{2\pi} |g(t, y)| |x_y| dt \\ &\quad + \int_0^{2\pi} |f(t, 0)| dt. \end{aligned} \quad (15)$$

Using (12), (14), and (15), there is constant $E > 0$, such that

$$|x_y^{(2n)}(t)| \leq EL, \quad \text{for all } t \in [0, 2\pi],$$

which, with (12) and (14), completes the proof of Lemma 2.

THEOREM C. Assume that $h(t) = h(t + 2\pi)$ is continuous and satisfies

$$m \leq |h(t)| \leq M \quad \text{for all } t \in [0, 2\pi],$$

where m, M are constants, and $c_i, i = 0, 1, \dots, n-1$, satisfies the inequality (4). Then the equation

$$x^{(2n+1)} + \sum_{i=0}^{n-1} c_i x^{(2i+1)} + h(t)x = 0 \quad (16)$$

has only a zero solution.

The proof of Theorem C is similar to that of Lemma 2. We only need to note that in (12), (14), and (15), $L = 0$,

THEOREM D. Assume that $h(t)$, c_i , $i = 0, 1, \dots, n - 1$, satisfies the conditions in Theorem C. Then the linear differential equation

$$x^{(2n+1)} + \sum_{i=0}^{n-1} c_i x^{(2i+1)} + h(t)x = e(t), \quad (17)$$

where $e(t) = e(t + 2\pi)$ is continuous, has a unique 2π -periodic solution.

Proof. By Theorem C, (16) has only a zero solution. Since the equation (17) is linear, the uniqueness implies the existence. The proof is completed.

Proof of Theorem B. First prove the uniqueness. Let $x_1(t)$ and $x_2(t)$ be any two 2π -periodic solution of (1). Then $x(t) = x_1(t) - x_2(t)$ is a 2π -periodic solution of the following equation:

$$x^{(2n+1)} + \sum_{i=0}^{n-1} c_i x^{(2i+1)} + \int_0^1 f_x(t, x_2(t) + \theta x(t)) d\theta x = 0.$$

From the hypothesis (H1), we see

$$m \leq \left| \int_0^1 f_x(t, x_2(t) + \theta x(t)) d\theta \right| \leq M.$$

Hence by Theorem C, $x(t) \equiv 0$, on R .

We next prove the existence. For each $y \in X$, define

$$Py = x_y,$$

where x_y is a 2π -periodic solution of $(6)_y$. By Theorem D, P is well defined.

Let $x_n, x \in X$ such that $\|x_n - x\| \rightarrow 0$, as $n \rightarrow \infty$. We assert: $\|Px_n - Px\| \rightarrow 0 (n \rightarrow \infty)$. If not, by Lemma 2, $\{Px_n\}$ would have a convergent subsequence, still denoting it by $\{Px_n\}$, such that $\|Px_n - y\| \rightarrow 0 (n \rightarrow \infty)$ and $Px \neq y$. Set $u_n = Px_n - Px$. Then $u_n(t)$ satisfies

$$u_n^{(2n+1)} + \sum_{i=0}^{n-1} c_i u_n^{(2i+1)} + g(t, x_n)u_n + (g(t, x_n) - g(t, x))Px = 0,$$

and u_n is 2π -periodic. Hence by the continuity of g , as $n \rightarrow \infty$, we get

$$u_0^{(2n+1)} + \sum_{i=0}^{n-1} c_i u_0^{(2i+1)} + g(t, x)u_0 = 0, \quad (18)$$

where $u_0(t) = y(t) - Px(t)$. By Theorem C, (18) has only the zero solution, which contradicts $y \neq Px$. By (18) and the Arzela–Ascoli theorem, it follows that P is completely continuous. From Lemma 2, PX is bounded. Applying the Schauder fixed-point theorem, P has a fixed point $x(t)$ in X , which is a solution of (1). This completes the proof of Theorem B.

Now we prove Theorem A by applying the continuation theorem of Mawhin (see, e.g., [5] or [6]).

Consider the following auxiliary equations:

$$x^{(2n+1)} + \sum_{i=0}^{n-1} c_i x^{(2i+1)} + \lambda g(t, x)x = -\lambda f(t, 0), \quad 0 < \lambda \leq 1, \quad (19)_\lambda$$

using the notation in (5). Define the linear operator $L: C_{2\pi}^{2n} \rightarrow L^2(0, 2\pi)$ by

$$Lx = x^{(2n+1)} + \sum_{i=0}^{n-1} c_i x^{(2i+1)}, \quad x \in C_{2\pi}^{2n}.$$

By (2) in Condition (H1), the operator $x(\cdot) \rightarrow f(\cdot, x(\cdot))$ maps L^2 into itself.

Multiplying each side of $(19)_\lambda$ by $x(t)$, integrating over $[0, 2\pi]$, and using the boundary conditions, we get

$$\lambda \int_0^{2\pi} g(t, x(t)) x^2(t) dt = -\lambda \int_0^{2\pi} f(t, 0) x(t) dt.$$

Dividing out λ and using $|g(t, x)| \geq m$ for all (t, x) , we get

$$\int_0^{2\pi} x^2(t) dt \leq M_0 \quad (20)$$

for a constant M_0 independent of x and λ . This gives an *a priori* bound in L^2 .

Instead of working in L^2 , one could work in $C_{2\pi}$, using (20) and the equation $(19)_\lambda$ to obtain a bound in $C_{2\pi}$.

Clearly, according to (H1) and the definition of topological degree, the mapping from the constants (the kernel of the operator L) to the constants, given by

$$\bar{x} \rightarrow \frac{1}{2\pi} \int_0^{2\pi} f(t, \bar{x}) dt, \quad \bar{x} \in R,$$

has nonzero topological degree on a large interval containing 0.

Employing the continuation theorem of Mawhin, we can complete the proof of Theorem A.

ACKNOWLEDGMENTS

The authors thank the referee for valuable advice.

REFERENCES

1. A. C. Lazer, Application of a lemma on bilinear forms to a problem in nonlinear oscillations, *Proc. Amer. Math. Soc.* **33**, No. 1 (1972), 89–94.
2. K. J. Brown and S. S. Lin, Periodically perturbed conservative systems and a global inverse function theorem, *Nonlinear Anal.* **4** (1979), 193–201.
3. Cong Fuzhong, Periodic solutions for $2k$ th order ordinary differential equations with nonresonance, *Nonlinear Anal.* **32**, No. 6 (1997), 787–793.
4. Li Yong and Wang Huaizhong, Periodic solutions of high order Duffing equations, *Appl. Math. J. Chinese Univ.* **6** (1991), 407–412.
5. R. E. Gaines and J. L. Mawhin, Coincidence degree and nonlinear differential equations, *Lecture Notes in Math.* **568** (1977).
6. J. L. Mawhin, “Topological Degree Methods in Nonlinear Boundary Value Problems,” CBMS 40, Amer. Math. Soc., Providence, 1980.